

New J/ψ suppression data and the comovers interaction model

A. Capella^{a)} and D. Sousa^{b)}

^{a)} Laboratoire de Physique Théorique¹

Université de Paris XI, Bâtiment 210, F-91405 Orsay Cedex, France

^{b)} ECT*, Villa Tambosi

Strada delle Tabarelle 286, 38050 Villazzano (Trento), Italy

Abstract

New data on the J/ψ suppression both in proton-nucleus and in lead-lead interactions have been presented recently by the NA50 collaboration. We show that these data, together with the final ones on sulfur-uranium interactions, can be described in the framework of the comovers interaction model with a unique set of three parameters : the nuclear absorption cross-section, the comovers interaction cross-section and a single (rescaled) absolute normalization. Expectations for J/ψ suppression at RHIC are also discussed.

LPT Orsay 03-17

March 2003

¹Unité Mixte de Recherche UMR n° 8627 - CNRS

1 Introduction

Before the Quark Matter conference of 2002, the NA50 interpretation of the data on J/ψ suppression was as follows [1-3]. The pA , SU and peripheral $Pb Pb$ data (up to $E_T \sim 35 \div 40$ GeV) can be described with nuclear absorption alone, with an absorptive cross-section $\sigma_{abs} = 6.4 \pm 0.8$ mb. At $E_T \sim 40$ GeV there is a sudden onset of anomalous suppression, followed by a steady fall off at larger E_T . However, at variance with this view, the most peripheral points in $Pb Pb$ collisions lied above the NA50 nuclear absorption curve – which extrapolates pA and SU data.

Two important sets of new data have been presented recently [4] [5]. The new NA50 data on pA reactions at 450 GeV/c indicate a smaller value of σ_{abs} than the one given above. However, within errors, pA and SU data can still be described with a single value of the absorptive cross-section $\sigma_{abs} = 4.4 \pm 0.5$ mb – substantially lower than the previous one [5]. The new, preliminary, $Pb Pb$ data [4], taken in 2000 with a target under vacuum, are consistent with previous ones except for the most peripheral ones – which are now lower and consistent with the nuclear absorption curve [4]. In this way, the NA50 interpretation remains valid. However, the new data lend support to the interpretation based on comovers interaction – according to which some anomalous suppression is already present in SU collisions. Indeed, in the comovers approach² the sudden onset of anomalous suppression due to deconfinement is replaced by a smooth anomalous suppression due to comovers interaction. The effect of the comovers turns out to be negligibly small in pA but it is sizable in SU interactions. With the smaller value of σ_{abs} from the new pA data, there is more room for comovers in SU .

The purpose of this work is to study the consistency of the new data on pA and $Pb Pb$ interactions, together with the final SU ones, with the comovers interaction model [8] [9]. We proceed as follows. Since the effect of the comovers suppression is sizable in SU , but negligibly small in pA , σ_{abs} has to be determined from the pA data alone. In previous works [8-9] we have used a value $\sigma_{abs} = 4.5$ mb, as a compromise between NA38/NA51 [10] and E537 [11] pA data. Actually, it has been shown in [12] that the old pA data are also consistent with $\sigma_{abs} = 4.5$ mb. As mentioned above,

²For reviews on deconfining and comover interaction models see [6]. Alternative models have also been proposed [7].

this value has now been confirmed by the recent NA50 data [5], and will be used throughout this paper. The second parameter in the model, the comovers interaction cross-section σ_{co} , can then be determined from the centrality dependence of the J/ψ suppression in $Pb\ Pb$ collisions. Obviously its value is correlated with that of the third parameter of the model, the absolute normalization. This normalization, in turn, is strictly related to the one in SU . The ratio of the $Pb\ Pb$ to the SU normalizations is equal to 1.051 ± 0.026 [5]. This is a rescaling factor which takes into account both isospin and energy corrections. In this way, with σ_{abs} fixed, the model is strongly constrained.

The main drawback of the comovers model [8-9] was precisely a mismatch of about 30 % between the absolute normalizations in SU and $Pb\ Pb$ [12]. The origin of this mismatch is the following. The high values of the most peripheral $Pb\ Pb$ data in the former NA50 analysis required a value $\sigma_{co} = 1$ mb. As stated above, in the new data the most peripheral $Pb\ Pb$ points are substantially lower and require a lower value of $\sigma_{co} = 0.65$ mb. Indeed, a smaller value of σ_{co} (with σ_{abs} fixed) leads to a flatter centrality dependence of the J/ψ suppression. This change in σ_{co} induces a change in the absolute normalization – which is now in good agreement with the (rescaled) one obtained in SU .

The plan of this paper is as follows. In Section 2 we present a short summary of the comovers interaction model [8-9]. In Section 3 we apply it to $Pb\ Pb$ collisions, where the data allow an accurate determination of both the comovers cross-section and the absolute normalization. We also compute the correlation between E_T and E_{ZDC} – the energy of the zero degree calorimeter and discuss the J/ψ suppression in the E_{ZDC} analysis. In Section 4 we show that the pp , pA and SU data can be described using the same values of σ_{abs} and σ_{co} as in $Pb\ Pb$ and a single (rescaled) normalization – obtained from either SU or $Pb\ Pb$ data. Section 5 contains our conclusions and expectations for J/ψ suppression at RHIC.

2 Comovers interaction in the dual parton model

The cross-section of minimum bias (MB), lepton pair (DY) and J/ψ event

samples are given by

$$I_{MB}^{AB}(b) \propto \sigma_{AB}(b) \quad (1)$$

$$I_{AB}^{DY}(b) \propto \int d^2s \sigma_{AB}(b) n(b, s) \quad (2)$$

$$I_{AB}^{J/\psi}(b) \propto \int d^2s \sigma_{AB}(b) n(b, s) S_{abs}(b, s) S_{co}(b, s) . \quad (3)$$

Here $\sigma_{AB}(b) = \{1 - \exp[-\sigma_{pp} AB T_{AB}(b)]\}$ where $T_{AB}(b) = \int d^2s T_A(s) T_B(b-s)$, and $T_A(b)$ are profile functions obtained from the Woods-Saxon nuclear densities [13]. Upon integration over b we obtain the AB total cross-section, σ_{AB} . $n(b, s)$ is given by

$$n(b, s) = AB \sigma_{pp} T_A(s) T_B(b-s) / \sigma_{AB}(b) . \quad (4)$$

Upon integration over s we obtain the average number of binary collisions $n(b) = AB \sigma_{pp} T_{AB}(b) / \sigma_{AB}(b)$.

The factors S_{abs} and S_{co} in (3) are the survival probabilities of the J/ψ due to nuclear absorption and comovers interaction, respectively. They are given by [8] [9]

$$S^{abs}(b, s) = \frac{[1 - \exp(-AT_A(s) \sigma_{abs})][1 - \exp(-B T_B(b-s) \sigma_{abs})]}{\sigma_{abs}^2 AB T_A(s) T_B(b-s)} \quad (5)$$

$$S^{co}(b, s) = \exp \left[-\sigma_{co} \frac{3}{2} N_{y_{DT}}^{co}(b, s) \ell n \left(\frac{\frac{3}{2} N_{y_{DT}}^{co}(b, s)}{N_f} \right) \right] \quad (6)$$

In (6), $N_{y_{DT}}^{co}(b, s)$ is the density of charged comovers (positives and negatives) in the rapidity region of the dimuon trigger and $N_f = (3/\pi R_p^2)(dN/dy)_{y^*=0} = 1.15 \text{ fm}^{-2}$ [8, 9, 14] is the corresponding density in pp . The factor $3/2$ in (6) takes care of the neutrals. In the numerical calculations we use $\sigma_{abs} = 4.5 \text{ mb}$. The value of σ_{co} and the absolute normalization will be determined from the data.

In order to compute the density of comovers we use the DPM formalism described in [15]. It turns out that the density of charged particles is given by a linear superposition of the density of participants and the density of binary collisions with coefficients calculable in DPM. All details can be found in [8] and [15].

Eqs. (1) to (6) allow to compute the impact parameter distributions of the MB , DY and J/ψ event samples. Experimental results are plotted as a function of

observable quantities such as E_T – the energy of neutrals deposited in the calorimeter. Using the proportionality between E_T and multiplicity, we have

$$E_T(b) = \frac{1}{2} q N_{y_{cal}}^{co}(b) \quad . \quad (7)$$

Here the multiplicity of comovers is determined in the rapidity region of the E_T calorimeter. The factor 1/2 is introduced because N^{co} is the charged multiplicity whereas E_T refers to neutrals. In this way q is close to the average transverse energy per particle, but it also depends on the calibration of the calorimeter. The correlation $E_T - b$ is parametrized in the form [3, 14]

$$P(E_T, b) = \frac{1}{\sqrt{2\pi qaE_T(b)}} \exp \left\{ -(E_T - E_T(b))^2 / 2qaE_T(b) \right\} \quad . \quad (8)$$

The E_T distributions of MB , DY and J/ψ are then obtained by folding Eqs. (1)-(3) with $P(E_T, b)$, i.e.

$$I_{AB}^{MB,DY,J/\psi}(E_T) = \int d^2b I_{AB}^{MB,DY,J/\psi}(b) P(E_T, b) \quad . \quad (9)$$

The parameters q and a are obtained from a fit of the E_T distribution of the MB event sample³. Note that since $N_{y_{cal}}^{co}(b)$ is nearly proportional to the number of participants (see Fig. 1 of [8]), our fit is practically identical to the one obtained [14] using the wounded nucleon model. Actually, we obtain identical curves to the ones in Fig. 1 of ref. [3] – where the E_T distributions of MB events of 1996 and 1998 are compared with each other. The values of the parameters for the 1996 data are $q = 0.62$ GeV and $a = 0.825$. For the 1998 data, the tail of the E_T distribution is steeper, and we get $q = 0.62$ GeV and $a = 0.60$ ⁴. In the following we shall use

³Note that the same value of the parameter a is used in the MB , DY and J/ψ event sample. A priori there could be some differences in the fluctuations for hard and soft processes. Actually, it has been claimed in Ref. [8] that there is a small shift in E_T between minimum bias, on one hand, and J/ψ or Drell-Yan pair production on the other hand – induced by the dimuon trigger. However, this is of no relevance for the present work, since, so far, the only 2000 data available have been obtained in the so-called standard analysis – in which the genuine ratio of J/ψ and DY cross-sections is measured.

⁴At first sight these sets of values look very different from the ones used by the NA50 collaboration. Nevertheless, they reproduce the same E_T distribution. This is due to the fact that the product qa , which according to Eq. (8), determines the width of the distribution, is very similar in the two cases. As for the difference in the values of q it is just due to its definition, which is different in the two approaches (Eq. (7), in our case).

the latter values. Indeed, according to the NA50 collaboration [2], the 1996 data (thick target) at large E_T are contaminated by rescattering effects – and only the 1998 data should be used beyond the knee.

The model described above allows to compute the E_T distribution of MB , DY and J/ψ event samples between peripheral AB collisions and the knee of the E_T distribution. Beyond it, most models, based on either deconfinement or comovers interaction, give a ratio of J/ψ to DY cross-sections which is practically constant – in disagreement with NA50 data. A possible way out was suggested in [9]. The idea is that, since E_T increases beyond the knee due to fluctuations, one can expect that this is also the case for the density of comovers. Since $N_{y_{DT}}^{co}$ does not contain this fluctuation, it has been proposed in [9] to introduce the following replacement in Eq. (6) :

$$N_{y_{DT}}^{co}(b, s) \rightarrow N_{y_{DT}}^{Fco}(b, s) = N_{y_{DT}}^{co}(b, s) F(b) \quad (10)$$

where $F(b) = E_T/E_T(b)$. Here E_T is the measured value of the transverse energy and $E_T(b)$ is its average value given by Eq. (7) – which does not contain the fluctuations.

3 J/ψ suppression in Pb Pb

a) E_T analysis

The new data [4] for the ratio of J/ψ over DY cross-sections versus the energy of the E_T calorimeter are shown in Fig. 1. They are compared with the results of the comovers interaction model described in Section 2. As explained there, there are two free parameters in the model ($\sigma_{abs} = 4.5$ mb has been fixed) : the comovers interaction cross-section σ_{co} (which controls the centrality dependence of the ratio) and the absolute normalization. A good description of the data is obtained using $\sigma_{co} = 0.65$ mb and an absolute normalization of 47.

The only difference between our result and the one in [8] resides in the value of σ_{co} . Since the effect of the comovers increases with centrality, a larger (smaller) value of σ_{co} leads to a larger (smaller) variation of the ratio of J/ψ over DY cross-sections between peripheral and central collisions. As mentioned in the Introduction, in the new NA50 analysis the values of this ratio for peripheral collisions are smaller. In

order to describe the new data, the value of σ_{co} has to be reduced. The curve in Fig. 1 corresponds to a reduction of σ_{co} from 1 mb (used in [8]) to 0.65 mb.

Since the values of σ_{co} and of the absolute normalization are correlated, the decrease of σ_{co} induces, in turn, a decrease of the absolute normalization. While in [8] the value of the absolute normalization was about 30 % higher [12] than in *SU*, the one in Fig. 1 is in good agreement with the *SU* one. This will be shown in the next section. It is interesting that almost the same value of σ_{co} ($\sigma_{co} = 0.7$ mb) was obtained in [16] from an analysis of *SU* data and former *Pb Pb* data [1] which covered a much smaller centrality range. In [16] the absolute normalizations in *SU* and *Pb Pb* were in good agreement with each other.

The *DY* cross-section in Fig. 1 has been integrated in the dimuon mass range 2.9 to 4.5 GeV. Since the J/ψ peak is inside this range, a model is needed in order to determine the *DY* cross-section. In the *SU* analysis, the GRV parton distribution functions at leading order (LO) have been used. Therefore, in order to use the (rescaled) absolute normalization of the *SU* data in *Pb Pb* (or vice versa), the same GRVLO distributions have to be used in the latter. This is the case in Fig. 1. In *Pb Pb* collisions, NA50 has also analyzed the data using, instead, MRS 43 distributions. They have found [4] that in this case the absolute normalization is lower by about 10 %. The comparison of the comovers model with the data is presented in Fig. 2. The absolute normalization is 43. The values of σ_{abs} and σ_{co} are, of course, unchanged.

b) E_{ZDC} analysis

The NA50 collaboration has also measured the J/ψ suppression in *Pb Pb* as a function of the energy of the zero degree calorimeter (E_{ZDC}). In Fig. 1, the results of this analysis have been plotted as a function of E_T , using the measured correlation between average values of E_T and E_{ZDC} . We see from Fig. 1 that the data obtained in the two analysis are consistent with each other, even for very central events, beyond the knee of the E_T and E_{ZDC} distributions. This important result has been predicted in [12]. Its physical origin is the following.

The energy of the zero degree calorimeter is given by

$$E_{ZDC}(b) = [A - n_A(b)]E_{in} + \alpha n_A(b) E_{in} . \quad (11)$$

Here $n_A(b)$ is the average number of participants at fixed b :

$$n_A(b) = A \int d^2s T_A(s) [1 - \exp \{-\sigma_{pp} B T_B(b-s)\}] \sigma_{AB}(b) \quad (12)$$

$A - n_A(b)$ is the number of spectator nucleons of A and $E_{in} = 158$ GeV is the beam energy. While the first term in the r.h.s. of Eq. (11) gives the bulk of E_{ZDC} , the latter corresponds to the contamination by secondaries emitted very forward [17] – assumed to be proportional to the number of participants, $n_A(b)$. Here also the value of α can be precisely determined from the position of the “knee” of the E_{ZDC} distribution of the MB event sample measured by NA50 [17]. We obtain $\alpha = 0.076$ [12].

Eqs. (7) and (11) give the relation between b and E_T and b and E_{ZDC} , respectively. These relations refer to average values and do not contain any information about the tails of the E_T or E_{ZDC} distributions. Eqs. (7) and (11) also lead to a correlation between (average values of) E_T and E_{ZDC} . This correlation [12] gives a good description of the experimental one [3]. It is practically a straight line⁵ and therefore can be accurately extrapolated beyond the knee of the E_T and E_{ZDC} distributions. It turns out that this extrapolation describes the measured $E_T - E_{ZDC}$ correlation quite well⁶ – even for values of E_T and E_{ZDC} in the tails of the distributions. This result suggests a correlation between b and E_{ZDC} of the form

$$P(E_{ZDC}, b) = P(E_T, b) \delta(E_T - E_T(E_{ZDC})) . \quad (13)$$

Folding (1) and (13) we obtain the E_{ZDC} distribution of MB events. It describes [12] the one measured by NA50, not only up to the knee, but also in the tail of the distribution. This result shows that the J/ψ suppression versus E_{ZDC} is just obtained from the corresponding one versus E_T by applying the $E_{ZDC} - E_T$ correlation, even for very central events beyond the knee of the distributions. In the 1996 and 1998 NA50 data, the J/ψ suppression versus E_{ZDC} indicated some features (“snake shape”) not present in the ones versus E_T . Such differences are no longer present in the new data.

⁵This is due to the fact that $N_{yca}^{co}(b)$ in Eq. (7) is practically proportional to $n_A(b)$ (see Fig. 1 of [8]).

⁶One can understand the physical origin of this extrapolation if one assumes that a fluctuation in E_T is essentially due to a fluctuation in n_A – which, in turn, produces a corresponding fluctuation in E_{ZDC} , via Eq. (11).

4 J/ψ suppression in pA and SU

Let us compute next the ratio R of J/ψ over DY cross-sections in SU at 200 GeV/c per nucleon in our model. We use, of course, the same values of the parameters as in $Pb\ Pb$: $\sigma_{abs} = 4.5$ mb and $\sigma_{co} = 0.65$ mb. To get this ratio versus b , the only new ingredient is the multiplicity of comovers – which is again computed in DPM, in the way described in Section 2. To compute $R(E_T)$, we also need the $E_T - b$ correlation in SU , which is parametrized as in Eq. (8). The parameters q and a have been obtained from a fit of the E_T -distribution of DY given in [18]. We obtain $q = 0.69$ GeV and $a = 1.6$. ($R(E_{ZDC})$ has not been measured in SU). In SU , data do not extend beyond the knee of the E_T -distribution. Therefore, effects such as E_T fluctuations, Eq. (10), are not relevant here.

Our results are shown in Fig. 3. We see that the E_T dependence of the suppression is reproduced. This indicates that there is, indeed, room in SU for the (comparatively small) suppression by comovers. As discussed above this could also be inferred from the different central values of σ_{abs} obtained in pA and SU . The absolute normalization of the curve in Fig. 3 is 45. Thus the normalizations in $Pb\ Pb$ and SU are consistent with each other. This normalization is 4 % smaller than the one obtained from the $Pb\ Pb$ data – in perfect agreement with the rescaling factor discussed in the Introduction.

In pA collisions, the effect of the comovers is negligible and, therefore, the description of the pA data is the same as in the NA50 analysis [5], since, as discussed above, the value of σ_{abs} they obtain is practically identical to ours. The corresponding normalization is 20 % higher than the one we have obtained in SU . This is also consistent with the expected rescaling factor between the two systems – which takes into account the difference in energy as well as in the rapidity regions covered by the dimuon trigger.

5 Conclusions and outlook

The NA50 deconfining scenario has been described in the Introduction. In this work we have presented a different scenario in which the sudden onset of anomalous suppression due to deconfinement is replaced by a smooth one resulting from

comovers interaction. This anomalous suppression is already present in SU and peripheral $Pb\ Pb$ collisions.

We have presented a description of the NA38/NA50 data on the J/ψ suppression in pp , pA , SU and $Pb\ Pb$ interactions in a comovers model. The model is strongly constrained by the existing data. This can be seen in the following way. In SU , the effect of the comovers is rather small and, thus, once the value of $\sigma_{abs} = 4.5$ mb is fixed, the absolute normalization depends little on the exact value of σ_{co} . Since the normalizations in SU and $Pb\ Pb$ are strictly related, we are left with a single free parameter, σ_{co} , to determine the J/ψ suppression in $Pb\ Pb$ (both in absolute value and centrality dependence). The model is, thus, strongly constrained and provides a unified description of the data in the various systems. On the other hand, it is not possible to describe the former NA50 data on $Pb\ Pb$ collisions for peripheral events in a consistent way. Indeed, these data require a value $\sigma_{co} = 1$ mb. As shown in [12], this, in turn, leads to a mismatch of about 30 % between the absolute normalizations in SU and $Pb\ Pb$ systems. Furthermore, with equal normalizations in SU and in $Pb\ Pb$, the J/ψ suppression is always larger in the latter than in the former, even for very peripheral events (see Fig. 7 of [12]).

Let us discuss briefly the expectations for J/ψ suppression at RHIC in the comovers interaction model. The calculation of the survival probability S_{co} is quite safe. Indeed, since σ_{co} is a cross-section near threshold, the same value obtained at SPS should be used at RHIC. The situation is quite different for S_{abs} . Many authors assume that σ_{abs} is the same at RHIC and at SPS. It has also been suggested that it can be significantly larger at RHIC. However, it seems plausible that at mid-rapidities, nuclear absorption at RHIC is small due to the fact that, contrary to SPS, the $c\bar{c}$ pair is produced outside the colliding nuclei. It is therefore crucial to have data on J/ψ production in pA interactions at RHIC. If $S_{abs} \sim 1$, the J/ψ suppression at RHIC and SPS will be comparable, since the smallness of the nuclear absorption will be approximately compensated by the increase of the comovers suppression – due to a larger comovers density at RHIC. Very preliminary data tend to indicate that this is indeed the case. Detailed calculations will be presented elsewhere.

A quantitative analysis of the new NA50 data in the deconfining scenario is still missing. On the other hand, the centrality dependence of the average p_T of J/ψ is

better described in the comovers approach than in a deconfining scenario [19]. At RHIC energies, a small nuclear absorption in pA collisions (i.e. $S_{abs} \sim 1$), would be a very interesting situation in order to discriminate between the comovers interaction model and a deconfining scenario. Indeed, in the latter, the shape of the centrality dependence would be almost flat for peripheral collisions (below the deconfining threshold) and would decrease above the threshold. Such a behavior would be a clear signal of deconfinement. On the contrary, in the comovers scenario, the fall-off would be continuous, from peripheral to central collisions, and determined by the same value of σ_{co} obtained from CERN SPS data.

Acknowledgements

We thank N. Armesto for discussions and M. Gonin, L. Kluberg and E. Scomparin for information on the NA50 data.

References

- [1] NA50 collaboration, M. C. Abreu et al, Phys. Lett. **B410**, 327 (1997); ibid page 337.
- [2] NA50 collaboration, M. C. Abreu et al., Phys. Lett. **B450**, 456 (1999).
- [3] NA50 collaboration, M. C. Abreu et al., Phys. Lett. **B477**, 28 (2000).
- [4] NA50 collaboration, L. Ramello in Proceedings Quark Matter 2002, Nantes (France), 18-24 July 2002 (to be published).
- [5] NA50 collaboration, P. Cortese in Proceedings Quark Matter 2002, ibid.
NA50 collaboration, R. Sahoyan in Proceedings XXVII Rencontres de Moriond, Les Arcs (France), March 2002.
- [6] C. Gerschel and J. Hufner, Ann. Rev. Nucl. Part. Sci. **49**, 255 (1999).
R. Vogt, Phys. Rep. **310**, 197 (1999).
H. Satz, Rep. Prog. Phys. **63**, 1511 (2000).
- [7] J. Qiu, J. P. Vary, X. Zhang, hep-ph/9809442;
A. K. Chaudhuri, preprint Variable Energy Cyclotron Center, Calcutta, India (September 14, 2001).
- [8] A. Capella, A. Kaidalov and D. Sousa, Phys. Rev. **C65**, 054908 (2002).
- [9] A. Capella, E. G. Ferreira and A. Kaidalov, Phys. Rev. Lett. **85**, 2080 (2000).
N. Armesto et al., Nucl. Phys. **A698**, 583 (2002).
- [10] NA38 collaboration, M. C. Abreu et al, Phys. Lett. **B466**, 408 (1999);
NA51 collaboration, M. C. Abreu et al., Phys. Lett. **B438**, 35 (1998).
- [11] FNAL E537 collaboration, M. Leitch et al., Phys. Rev. Lett. **84**, 3256 (2000).
- [12] A. Capella and D. Sousa, nucl-th/0110072, unpublished.
- [13] C. W. Jager et al., Atomic Data and Nuclear Tables **14**, 485 (1974).
- [14] D. Kharzeev, C. Lourenço, M. Nardi and H. Satz, Z. Phys. **C74**, 307 (1997).

- [15] A. Capella and D. Sousa, Phys. Lett. **B511**, 185 (2001).
- [16] N. Armesto and A. Capella, Phys. Lett. **B430**, 23 (1998).
- [17] NA50 collaboration, M. C. Abreu et al., Phys. Lett. **B521**, 195 (2001).
- [18] NA38 collaboration, M. C. Abreu et al., Phys. Lett. **B449**, 128 (1999).
- [19] N. Armesto, A. Capella and E. G. Ferreira, Phys. Rev. **C59**, 395 (1999).
A. K. Chaudhuri, nucl-th/0212046.

Figure Captions

Fig. 1 : Ratio of J/ψ to DY cross-sections versus E_T in $Pb\ Pb$ collisions at 158 GeV/c per nucleon (solid line). The preliminary data are from [4]. GRVLO parton distribution functions have been used in order to calculate the DY cross-section in the mass range 2.9 to 4.5 GeV.

Fig. 2 : Same as in Fig. 1, using MRS 43 parton distribution functions.

Fig. 3 : The ratio of J/ψ to DY cross-sections as a function of E_T in SU collisions at 200 GeV/c per nucleon. The data are from [18].




Figure 1




Figure 2




Figure 3